

The Bombardment of the Earth During the Hadean and Early Archean Eras: A New Look at the Late Accretion

S. Marchi¹, W.F. Bottke¹, L.T. Elkins-Tanton², A. Morbidelli³, K. Wuennemann⁴, D. A. Kring⁵, M. Bierhaus⁴

1. Southwest Research Institute, Boulder, CO, United States; 2. Carnegie Institution for Science, Washington, DC, United States; 3. Observatoire de la Cote d'Azur, Nice, France; 4. Museum für Naturkunde, Berlin, Germany; 5. USRA - Lunar and Planetary Institute, Houston, TX, United States.

Our knowledge of the Earth during the Hadean and early Archean eons (ca 4.5-3.5 Ga) is very limited, mainly because few rocks older than 3.8 Ga have been found (e.g. Harrison 2009). Hadean-era zircons have allowed us to glean important insights into this era, but their data has led to considerably different evolution models for the evolution of the early Earth; some predict a hellish world dominated by a molten surface with a sporadic steam atmosphere (e.g. Pollack 1997), while others have predicted a tranquil, cool surface with stable oceans (e.g. Wilde et al 2001; Valley et al 2002). To understand whether either model (or both) could be right, we believe it is useful to quantitatively examine the post Moon-forming impact bombardment of the early Earth.

Over the last several years, through a combination of observations (e.g., Marchi et al 2012), theoretical models (e.g., Bottke et al 2012), and geochemical constraints from lunar rock (e.g. highly siderophile elements -HSE- abundances delivered to the Moon by impactors; the global number of lunar basins; the record of Archean-era impact spherule beds on Earth; Walker 2009; Neumann et al 2012), we have constructed a calibrated model of the early lunar impactor flux (Morbidelli et al 2012). Our results have now been extrapolated to the Earth, where they can make predictions about its post-Moon formation bombardment, the so called *late accretion*.

Using a Monte Carlo code to account for the stochastic nature of major impacts, and constraining our results by the estimated HSE abundances of Earth's mantle (that were presumably delivered by impactors; Walker 2009; Bottke et al. 2010), we find the following trends. In the first ~100-200 Myr after the formation of the Moon, which we assume was created ~4.5 Ga, the Earth was almost entirely resurfaced by impacts. This bombardment, which included numerous $D > 1000$ km diameter impactors, should have vigorously mixed the crust and upper mantle. Between ~4.1-4.3 Ga, the impactor flux steadily decreased; though an uptick near ~4.1 Ga caused by the so-called Late Heavy Bombardment should have delivered a new round of large impactors striking at a mean velocity ~1.5 times higher than in previous epochs. Overall, only a relatively small fraction of ancient terrain survives unscathed all the way to the early Archean. We speculate that if impacts are involved with Hadean zircon formation, a scenario we find plausible, the apparent preference for ~4.1 Ga ages among Hadean zircons may be a combination of (i) terrain (and zircon) preservation and (ii) the ability of large impactors to create zircons (i.e., ~4.1 Ga would potentially be the "sweet spot" in this competition).

Bottke W.F., et al., Nature 485, 78, 2012.

Harrison T.M., Annu. Rev. Earth. Planet. Sci., 37, 2009.

Marchi S., et al., EPSL, 325, 2012.

Morbidelli A., et al., EPSL, 355, 2012.

Pollack H.N. Greenstone belts, Oxford University Press, 1997.

Valley J.W., et al., Geology, 30, 2002.

Walker R.J., Chem. Der Erd., 69, 2009.

Neumann G. et al., AGU 2012.